



International Journal of Public Health Science

Scopus coverage years: from 2020 to Present

Publisher: Intelektual Pustaka Media Utama

ISSN: 2252-8806 E-ISSN: 2620-4126

Subject area: Medicine: Medicine (miscellaneous) | Medicine: Health Policy | Social Sciences: Health (social science) |

Medicine: Public Health, Environmental and Occupational Health | Nursing: Nutrition and Dietetics

Source type: Journal

CiteScore 2021

0.4

SJR 2021

0.142

SNIP 2021

0.142

-
- [Statistical model for IC₅₀ determination of acetylcholinesterase enzyme for Alzheimer's disease](#) [PDF](#)
Anwar Fitrianto, Siau Man Mah, Siau Hui Mah 770-778
-
- [Hormonal contraceptive use related to breast cancer among women in Indonesia: a nationwide study](#) [PDF](#)
Solikhah Solikhah, Monthida Sangruangake, Ratu Matahari, Wulan Rahmadhani, Rochana Ruliyandari 779-784
-
- [Psychometric properties of the smartphone addiction proneness scale in a sample of Malaysian adolescents](#) [PDF](#)
Sii Jiing Chan, Kee Jiar Yeo, Lina Handayani 785-792
-
- [Behaviour of cervical cancer prevention among fertile age woman: health promotion approach](#) [PDF](#)
Julinda Malehere, Ni Ketut Alit Armini, Elida Ulfiana, Kartika Indaswari Dewi 793-799
-
- [Impact of VUCA world on children's emotional development during online learning](#) [PDF](#)
Afiz Azri Mohd Ghani, Lim Seong Pek, Rita Wong Mee Mee, Md Rosli Ismail, Uzzairah Nabila Ahmad Tazli, Tengku Shahrom Tengku Shahdan, Fatin Syamilah Che Yoh 800-807
-
- [The determinants of adolescent smokers in Indonesia](#) [PDF](#)
Sindu Setia Lucia, Novin Yetiani, Linda Suwarni, Heni Rusmitasari, Maretalinia Maretalinia, Suyitno Suyitno 808-814
-
- [Development and validation of a sexual and reproductive health knowledge scale](#) [PDF](#)
Shih-Hui Lee, Kee-Jiar Yeo, Lina Handayani 815-822
-

International Journal of Public Health Science (IJPHS)

Vol. 11, No. 3, September 2022, pp. 808~814

Editorial Team

Advisory Board

[Prof. Hans-Olov Adami](#), Harvard School of Public Health, United States
[Assoc. Prof. Dr. Luoping Zhang](#), University of California, Berkeley, United States
[Dr. Khalid M. Al Aboud](#), King Faisal Specialist Hospital and Research Centre, Saudi Arabia

Editor-in-Chief

[Prof. Dr. Jay G. Silverman](#), University of California, San Diego School of Medicine, United States

Managing Editor

[Dr. Lina Handayani](#), Universitas Ahmad Dahlan, Indonesia

Associate Editors

[Assoc. Prof. Dr. Rob Martinus Van Dam](#), National University of Singapore, Singapore
[Assoc. Prof. Dr. Mohd Hasni Jaafar](#), Universiti Kebangsaan Malaysia, Malaysia
[Assoc. Prof. Dr. Henry Odhiambo Imhonde](#), Ambrose Alli University, Nigeria
[Prof. Dr. Mane Abhay Babruwahan](#), Smt. Kashibai Navale Medical College, India
[Dr. Fazal Shirazi](#), MD Anderson Cancer Center, United States
[Dr. Miguel A. Mayer MD, PhD, MPH, MSc, Dipl](#), Universitat Pompeu Fabra, Spain
[Dr. Muhiuddin Haider](#), University of Maryland, United States
[Dr. Tassanee Rawiworakul](#), Mahidol University, Thailand
[Dr. Xiaoli Gao](#), University of Texas Health Science Center at San Antonio, United States
[Dr. Yuming Guo](#), Monash University, Australia

Editorial Board Members

[Prof. Dr. George Williams Rutherford](#), Global Health Sciences, University of California, San Francisco, United States
[Prof. Dr. Jens Aggaard-Hansen](#), Danish Bilharziasis Laboratory, Denmark
[Prof. Dr. Louise H. Taylor](#), Global Alliance for Rabies Control, Manhattan, United States
[Prof. Dr. Mogens Vyberg](#), Aalborg University, Denmark
[Dr. Abhinand Thavalappil](#), Ryerson University, Canada
[Dr. Stephanie S. Rothenberg](#), University of Pittsburgh Medical Center, United States
[Dr. Aletha Yvette S. Akers](#), Magee-Womens Hospital, Pittsburgh, United States
[Dr. Andrew Winokur](#), University of Connecticut School of Medicine, United States
[Dr. Barbara Abrams](#), University of California, United States
[Dr. Barbara K. Campbell](#), University of California, San Francisco, United States
[Dr. Christopher Bowie](#), University of Canterbury, New Zealand
[Dr. Erika Villavicencio-Ayub](#), Universidad Nacional Autónoma de México, Mexico
[Dr. Phumulani Mavis Mulaudzi](#), University of Pretoria, South Africa
[Dr. Jennifer A. Fish](#), The University of South Australia, Australia
[Dr. Li-Ping Zou](#), Department of Paediatrics, Chinese PLA General Hospital, China
[Dr. Phayong Thepakorn](#), Sirindhorn College of Public Health, Thailand
[Dr. Rob M.G. van Bommel](#), Catharina Hospital, Netherlands
[Dr. Tarik Beraket](#), University of Toronto, Canada
[Dr. Vera Fernandes](#), Hospital de Braga, Portugal

International Journal of Public Health Science (IJPHS)

p-ISSN: 2252-8806, e-ISSN: 2620-4126



The determinants of adolescent smokers in Indonesia

Sindu Setia Lucia¹, Novin Yetiani², Linda Suwarni³, Heni Rusmitasari⁴, Maretalinia Maretalinia^{5,6},
Suyitno Suyitno⁷

¹Department of Public Health, Faculty of Health Sciences, Universitas Respati Yogyakarta, Yogyakarta, Indonesia

²Department of Midwifery, Sekolah Tinggi Ilmu Kesehatan Kapuas Raya, Sintang, Indonesia

³Department of Public Health, Faculty of Health Sciences, Universitas Muhammadiyah Pontianak, Pontianak, Indonesia

⁴Department of Public Health, Faculty of Public Health, Universitas Muhammadiyah Semarang, Semarang, Indonesia

⁵Program of Demography, Institute for Population and Social Research, Mahidol University, Nakhon Pathom, Thailand

⁶Department of Public Health, Universitas MH Thamrin, Jakarta, Indonesia

⁷Department of Public Health, Institut Kesehatan Teknologi Graha Medika, Kotamobagu, Indonesia

Article Info

Article history:

Received Nov 4, 2021

Revised Apr 25, 2022

Accepted Jun 20, 2022

Keywords:

Adolescent

Cigarettes logo

Indonesia

Pocket money

Policy

Purchase near the school

ABSTRACT

In Indonesia, the prevalence of adolescence smokers is high. The situation might be worse during the epidemiological transition. This study aimed to estimate the sociodemographic factors related to tobacco use (light, moderate, and heavy) among adolescent in school age in Indonesia. This cross-sectional study used the raw data from the secondary data of Indonesia Global Youth of Tobacco Survey (GYTS), 2019. This study only included the 9,992 adolescent smokers in school age as the sample. School was selected based on the proportional probability based on the number of students. The class was selected by random sampling method and the students in that class were eligible to join the survey. This current study only selected those who smoked at the time of survey. The univariate, bivariate (Chi-square and t-test), and multivariate (ordinal logistic regression) has been tested in this study. The prevalence of tobacco uses among adolescent in Indonesia in 2019 was 19.2%. The determinants of tobacco use mostly related to pocket money, having products with cigarettes logo, and can purchase near the school. The role of sociodemographic, factors related to current tobacco use is very important to arrange the policy. The policy is very important to prevent and control tobacco use, especially among adolescence.

This is an open access article under the [CC BY-SA](#) license.



Corresponding Author:

Maretalinia Maretalinia

Institute for Population and Social Research, Mahidol University

Phutthamonthon Sai 4 Rd., Salaya, Phutthamonthon, Nakhon Pathom, Thailand

Email: maretalinia.21@gmail.com

1. INTRODUCTION

The smoking trend among adolescent is an old issues in developing countries, including Indonesia. The trend is increasing since the cigarette price in Indonesia is very cheap which was around 20,830 Indonesian rupiah (IDR) in average in 2020 [1]. The affordability of Indonesian to purchase the cigarette is still high even though the government has been applied the tax [2]. Applying the tax actually does not affect anything because the price only increased 1% [2]. As for student in adolescent age group, the money pocket is the main source to purchase the cigarette. However, the student could not find the cigarettes shop/retail near the school. The regulation from Indonesian Ministry of Education mentioned that the school environment is a free smoke area that need to be supported by headmaster of school, teachers, students, and others people in school area [3].

Smoking behavior is one unhealthy behavior as risk factor to non-communicable diseases (NCD) that may lead the hypertension, heart disease, stroke, and cancer [4]. The changing of phenomena from communicable diseases to non-communicable diseases will put the smokers as high-risk population. The condition will be worse if the students are also using the drug. The younger age of first try of cigarette, the higher risk to have noncommunicable diseases (NCD) [5]. Around 1.2 million passive smokers were estimated death due to the smoke exposed based on the data from World Health Organization (WHO) and seven million people deaths because their direct smoking behavior where Asia is the area with the highest number of smokers in the world [6].

Focusing on the adolescent in the school age, the risk factors of smoking is the complex one. The closest environment is in the family where adolescent spent much more time beside the school, and community. The study in Sumatera and Java Island found the fact that family and friends are the most influencer of early age smokers [7]–[9]. The tendency of children to imitate the parents is the nature condition that showed the effect of family role and environmental [10]. Another study using Indonesia Global Youth Tobacco Survey (GYTS) 2014 revealed that friends, parents, and teachers are the role mode to influence student to be a smoker [11].

GYTS is a global project managed by World Health Organization (WHO). The Indonesia GYTS is a survey designed to estimate the worldwide burden of tobacco use among adolescent [12]. In Indonesia, the tobacco users in adolescent school age always increase. It increased a lot from 0.3% in 1995 to 2.0% in 2007 [7], [8]. The data from basic health research Indonesia in 2013, prevalence of smokers among adolescent was 18.3% [13]. Comparing the place of residence, in 2017 smokers in rural (1.55%) is higher than urban (1.07%) [14]. The main findings from GYTS 2019 revealed that 19.2% students are using the tobacco but around 1% of them currently use the smokeless tobacco products [15].

Indonesia has been implementing the policies to control the tobacco use. The regulation from Minister of Education mentioned that the school environment is a free smoke area that need to be supported by headmaster of school, teachers, students, and others people in school area [15]. School is prohibited to produce, sell, distribute, sponsored, and put a cigarettes logo. The regulation about addictive product also mentioned that smoke cigarette is prohibited in enclosed public areas including school [16]. The regulation of Indonesian Ministry of Finance mentioned the price of cigarette stick only around IDR 1,350 (USD 0.094) which is cheap [17]. The cigarette advertisement is prohibited to demonstrate by using the real cigarette, and there is time restriction of promotion in television [18]. However, those policies are not implemented well because people can access to the cigarettes easily. For instance, adolescent still can purchase the cigarette without showing the identity card, which mean that the age restriction (18+) is not implement yet [19].

Study about determinants of smoking behavior is important. The effect of early smoke may associate with the adult smokers [20]. Most of adult smokers reported that they become addicted during adolescent [21]–[23]. The objective of this study was to examine the sociodemographic factors related to light, moderate, and heavy adolescent smokers based on the data from Indonesia Global Youth Tobacco Survey (GYTS) Indonesia, 2019.

2. RESEARCH METHOD

The GYTS is the cross-sectional study which has been conducted in 34 provinces in 2019. This survey aimed to control the tobacco use among adolescent in age 13 to 17 years [24]. In term of sampling method, school was selected based on the proportional probability. The class was selected by random sampling method and all of eligible student in grade 7-12 could join to the survey. Total sample in the original survey was 9,992 students. This study used the current tobacco use dependent variable based on the questions coded CR8: *Please think about the days you smoked cigarettes (including smoking white cigarettes, hand-rolled cigarettes or clove cigarettes) during the past 30 days. How many cigarettes did you usually smoke per day?* The sample was only the smokers who answered one or more days. After weighted to the place of residence and drop the missing data, total smokers in this study are 1,034 students.

The statistical analysis used in this study consists of univariate, bivariate, and multivariate. The univariate showed the frequency and percentage (for categorical variable) and minimum number, maximum number, and mean (for continuous variable) of each variable to see the characteristic of respondents. The bivariate analysis has been done by using Chi-square and t-test to see the crude odd ratio (COR) for each independent variable to dependent variables. The ordinal logistic regression has been done to examine all of independent variables to dependent variable in three models which was presented by adjusted odd ratio (AOR). STATA version 15 was used to analyze the data.

3. RESULTS AND DISCUSSION

3.1. Results

Totally, 19.2% adolescents were currently tobacco use with male percentage 35.6% and female 3.5%. In detail, Table 1 describes the characteristic of tobacco users. Mean age of adolescent smokers was 14 years old, the majority of them were male, and they mostly in grade 9 (junior high school). In terms of weekly pocket money, more than a half of them (60.74%) got less than IDR 30,000 (2.09 USD). The majority of them also did not have any products with cigarette logo (82.40%). Around 70% of adolescent smokers reported cigarettes with brand GG Mild and others brand as the brand usually smoke. Almost a half of them could not purchase cigarettes near school (47.20%).

Table 1. The characteristics of tobacco users

Variables (n=1,034)	Frequency	Percentage (%)
Age Min 11, Max 17, Mean 14.72		
Gender		
Female	51	4.93
Male	983	95.07
Grade Min 7, Max 12, Mean 9.40		
During an average week, how much money do you have that you can spend on yourself, however you want?		
Less than IDR 30,000 (USD 2.09)	628	60.74
More than IDR 30,000 (USD 2.09)	406	39.26
How much do you pay when you buy 1 cigarette?		
Less than Rp. 1,000 (0.07 USD)	231	22.34
IDR 1,000 (USD 0.07) – IDR 1,500 (USD 0.10)	566	54.74
IDR 1,600 (USD 0.11) – IDR 2,000 (USD 0.14)	171	16.54
More than IDR 2,000 (USD 0.14)	66	6.38
Do you have something (for example, t-shirt, pen, backpack, hat or sun glasses) with a cigarette product brand logo on it?		
No	852	82.40
Yes	182	17.60
During the past 30 days, what brand of cigarettes did you usually smoke?		
Unusual, A Mild, LA Lights	181	17.50
Mild Clas, Djarum Super MLD	122	11.80
GG Mild and others	731	70.70
Can you purchase cigarettes near your school?		
No	488	47.20
Yes	382	36.94
Do not know	164	15.86

Table 2 provides information about the relationship between each independent variable and dependent variable. In the sociodemographic variable, age, grade, money pocket, and had something with cigarette logo were statistically related to current tobacco use. The strongest independent variable in this domain is have something with logo.

Table 2. The bivariate results of current tobacco use

Variables (n=1,034)	Crude odd ratio	Confidence interval 95%	p-value
Sociodemographic			
Age	1.25	1.02-1.54	0.035
Gender	4.30	0.06-9.67	0.989
Grade	1.27	1.05-1.55	0.013
During an average week, how much money do you have that you can spend on yourself, however you want?	2.20	1.12-4.33	0.022
How much do you pay when you buy 1 cigarette?	0.93	0.61-1.42	0.748
Do you have something (for example, t-shirt, pen, backpack, hat or sun glasses) with a cigarette product brand logo on it?	3.52	1.78-6.96	<0.001
During the past 30 days, what brand of cigarettes did you usually smoke?	0.84	0.56-1.25	0.382
Can you purchase cigarettes near your school?	1.32	0.85-2.04	0.217

Note: the bold p-value means <0.05

Table 3 the multivariate results has been done to see the relationship and magnitude of independent variables and dependent variable. Those who had money pocket more than IDR 30,000 a week 2.32 times more likely to be heavy smokers compared by those had money pocket less than IDR 30,000. Having products with cigarette logo statistically associated with tobacco use, compare to those had no, those who had products with logo of cigarette 3.42 times more likely to be heavy smokers. In terms of accessibility to purchase cigarettes, those who could purchase near school were 2.75 times more likely to be heavy smokers.

Table 3. The ordinal logistic regression of current tobacco use

Variables (n= 1,034)	AOR	CI 95%
Age	1.02	0.72-1.45
Gender (ref: female)		
- Male	1.23	0.08-4.45
Grade	1.20	0.86-1.68
During an average week, how much money do you have that you can spend on yourself, however you want? (ref: Less than IDR 30,000)		
- More than or equal IDR 30,000	2.32*	1.13-4.74
How much do you pay when you buy one cigarette? (ref: Less than IDR 1,000)		
- IDR 1,000 – IDR 1,500	0.83	0.34-2.04
- IDR 1,600 – IDR 2,000	0.65	0.21-2.00
- More than IDR 2,000	0.32	0.04-2.72
Do you have something (for example, t-shirt, pen, backpack, hat or sun glasses) with a cigarette product brand logo on it? (ref: No)		
- Yes	3.42**	1.68-6.98
During the past 30 days, what brand of cigarettes did you usually smoke? (ref: Unusual, A Mild, LA Lights)		
- Mild Clas, Djarum Super MLD	0.54	0.14-2.12
- GG Mild and others	0.57	0.25-1.31
Can you purchase cigarettes near your school? (ref: No)		
- Yes	2.75*	1.24-6.09
- Do not know	1.28	0.38-4.22

Note: *p-value<0.05, **p-value<0.01, ***p-value<0.001

(Log likelihood = -157.31, LR chi2(12) =36.00, Prob >chi2 = 0.0003, Pseudo R2= 0.1027)

3.2. Discussion

The prevalence of current tobacco use among adolescent in Indonesia remains high compared with previous survey (in 2014) which 19.2% (35.6% male and 3.5% female) [24]. The prevalence of tobacco use in Indonesia was higher than Cameroon, Myanmar, Botswana, and India with current adolescent smoking was 11.2%, 13.6%, 10%, and 11.4% respectively [25]–[28]. Compared to 10 countries in Africa, that the current cigarettes smoking among students aged 13-15 was in range 3.4% to 13.6%, lower than Indonesia [29].

Sociodemographic and individual factors consisted of variables from the individual purchasing and indigenous. Higher money pocket is reflecting the high economic status and the high power of purchase the cigarette. In Indonesian context, students are given money pocket with daily or weekly frequencies. The money is using for meal, fulfill other school need. The logo of cigarettes on the something daily use may initiate people to try cigarette. The using of logo is also showing the identity of owner, including the identity as smokers. The access and affordability to purchase cigarettes near school may be due to the kiosk or shop near school sells the tobacco products. The knowledge of cigarette might come from the family who spend the majority of time with the adolescent [30].

The findings in this study emphasized the three main variables that determined the current tobacco use, such as money pocket, having products with logo, and can purchase near the school. These findings related to the policy and regulation in Indonesia. The regulation from Indonesian Ministry of Education mentioned that the school environment is a free smoke area that need to be supported by headmaster of school, teachers, students, and others people in school area [3]. School is prohibited to produce, sell, distribute, sponsored, and put a cigarettes logo. The regulation about addictive product also mentioned that smoke cigarette is prohibited in enclosed public areas including school [17]. The regulation of Indonesia Minister of Finance mentioned the price of cigarette stick [18]. The cigarette advertisement is prohibited to demonstrate by using the real cigarette, and there is time restriction of promotion in television [19], [31]–[33].

4. CONCLUSION

The prevalence of tobacco uses among adolescent in Indonesia in 2019 was 19.2%. The determinants of current tobacco use mostly related money pocket, to having products with cigarettes logo, and can purchase near the school. The role of sociodemographic, factors related to current tobacco use is very important to arrange the policy to prevent and control tobacco use.

The multilevel of prevention and control can reduce the prevalence of current tobacco use among adolescent, started from student level, school level, household level, community level, and national level. All of stakeholders need to take a part to prevent and control the prevalence of tobacco use. School and other stakeholder need to promote the harmfulness of smoking and smoke from other cigarettes, provide the health warning in the poster and describe in detail. Additionally, the parents are also need to control the money pocket so children will not use it to buy invaluable thing such as cigarettes. The government also need to control the policies and program that has been established and evaluate and compared it to other countries.





REFERENCES

- [1] Statista, "Average retail price of a packet of white cigarettes in Indonesia from 2011 to 2020," *Statista.com*, 2021.
- [2] B. W. Prasetyo and V. Adrison, "Cigarette prices in a complex cigarette tax system: Empirical evidence from Indonesia," *Tobacco Control*, vol. 29, no. 6, pp. 618–623, 2020, doi: 10.1136/tobaccocontrol-2018-054872.
- [3] Ministry of Education Indonesia, "The Regulation of Minister of Education of Indonesia Number 64 2015 concerning No Smoking Areas in School Environments stated that school principals, teachers, education staff, students, and other parties are prohibited from smoking," *Ministry of Education Republic of Indonesia*, 2015.
- [4] J. and M. K. J. Saari, A. J., Kentala, "The smoking habit of a close friend or family member - How deep is the impact? A cross-sectional study," *BMJ Open*, vol. 4, no. 2, pp. 1–6, 2014, doi: 10.1136/bmjopen-2013-003218.
- [5] M. P. Sylvestre *et al.*, "A tool to identify adolescents at risk of cigarette smoking initiation," *Pediatrics*, vol. 142, no. 5, 2018, doi: 10.1542/peds.2017-3701.
- [6] World Health Organization, "Tobacco - Key Facts," *Disponível em*: <https://www.who.int/news-room/fact-sheets/detail/tobacco>. Acesso em 27 Abril 2019., 2019.
- [7] T. Suryati and U. Tarigan, "Behavior tobacco consumption of junior high school student aged 13 – 15 years in Sumatra and Java (Analysis of Indonesia - Gyts 2009)," *Buletin Penelitian Sistem Kesehatan*, vol. 16, no. 3, pp. 259–266, 2013.
- [8] F. Jalilian *et al.*, "Socio-demographic characteristics associated with cigarettes smoking, drug abuse and alcohol drinking among male medical university students in Iran," *Journal of Research in Health Sciences*, vol. 15, no. 42–46, 2015.
- [9] M. Wibowo, L. Sofiana, S. M. Ayu, E. Gustina, and I. Khoeriyah, "Adolescent smoking behaviour determinants in the city of Yogyakarta, Indonesia," *International Journal of Community Medicine And Public Health*, vol. 6, no. 12, p. 5064, 2019, doi: 10.18203/2394-6040.ijcmph20195445.
- [10] I. J. Rickard, W. E. Frankenhuis, and D. Nettle, "Why are childhood family factors associated with timing of maturation? a role for internal prediction," *Perspectives on Psychological Science*, vol. 9, no. 1, pp. 3–15, 2014, doi: 10.1177/1745691613513467.
- [11] H. Jamal, A. Z. Abdullah, and M. T. Abdullah, "The Determinants of Social and Behavior of Tobacco Use Among Youth in Indonesia: An Analysis of Indonesia Global Youth Tobacco Survey in 2014" *Jurnal Kesehatan Vokasional*, vol. 5, no. 3, p. 141, 2020, doi: 10.22146/jkesvo.56718.
- [12] R. H. and M. J. Shafey O, Eriksen M, "The tobacco atlas, 4th edn. American Cancer Society," *Atlanta, GA, USA*, p. 30, 2010.
- [13] Ministry of Health Indonesia. (2019). The Result of Indonesian Basic Health Survey (Riset Kesehatan Dasar). <https://www.litbang.kemkes.go.id/laporan-riset-kesehatan-dasar-risikesdas/>
- [14] J.-H. Kim, J. Noh, J.-W. Choi, and E.-C. Park, "Association of education and smoking status on risk of diabetes mellitus: a population-based nationwide cross-sectional study," *International Journal of Environmental Research*, vol. 14, no. 6, p. 655, 2017.
- [15] Susenas, "Profile of Indonesian Children 2018, s.l.: Center for Statistics," *Jakarta: KKPPA (Kementerian Pemberdayaan Perempuan dan Perlindungan Anak)*, 2018.
- [16] Government Regulation of Indonesia Number 109 Year 2012 about Safeguarding of Materials Containing Addictive Substances in the Form of Tobacco Products for Health, President of Republic Indonesia 1 (2012). <https://peraturan.bpk.go.id/Home/Download/34279/PP%20Nomor%20109%20Tahun%202012.pdf>
- [17] The Regulation of Minister of Finance Indonesia No 198/PMK 010/2020, Pub. L. No. 198, Ministry of Finance of Republic of Indonesia 1 (2021). <https://jdih.kemenkeu.go.id/download/c52d2911-23cd-4d73-a017-ad9ffa5a0c3c/198~PMK.010~2020Per.pdf>
- [18] A. Lyons, A. McNeill, and J. Britton, "Tobacco imagery on prime time UK television," *Tobacco Control*, vol. 23, no. 3, pp. 257–263, 2014, doi: 10.1136/tobaccocontrol-2012-050650.
- [19] Syahrani, Prakoso, C. T., & Widyaningtyas, E. S. (2018). The Implementation of the Mayor's Regulation of a Non-Smoking Area (Study at the Smart Park of Samarinda City). *EJournal Administrasi Negara*, 6(1), 7117–7131.
- [20] World Health Organization. (2015). Adolescent Development. World Health Organization. http://www.who.int/maternal_child_adolescent/topics/adolescence/dev/en/. Accessed 27 Jan 2015
- [21] A. F. Buchmann *et al.*, "Early smoking onset may promise initial pleasurable sensations and later addiction," *Addiction Biology*, vol. 18, no. 6, pp. 947–954, 2013, doi: 10.1111/j.1369-1600.2011.00377.x.
- [22] B. A. Primack, S. Soneji, M. Stoolmiller, M. J. Fine, and J. D. Sargent, "Progression to traditional cigarette smoking after electronic cigarette use among us adolescents and young adults," *JAMA Pediatrics*, vol. 169, no. 11, pp. 1018–1023, 2015, doi: 10.1001/jamapediatrics.2015.1742.
- [23] S. Soneji *et al.*, "Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults a systematic review and meta-analysis," *JAMA Pediatrics*, vol. 171, no. 8, pp. 788–797, 2017, doi: 10.1001/jamapediatrics.2017.1488.
- [24] Ministry of Health Indonesia. (2019a). The Report of Indonesia Global Youth Tobacco Survey 2019. https://cdn.who.int/media/docs/default-source/searo/tobacco/global-youth-tobacco-survey/gyts-indonesia-extended-factsheet.pdf?sfvrsn=d202f34f_3
- [25] B. H. M. Ngahane, H. A. Ekobo, and C. Kuaban, "Prevalence and determinants of cigarette smoking among college students: A cross-sectional study in Douala, Cameroon," *Archives of Public Health*, vol. 73, no. 1, 2015, doi: 10.1186/s13690-015-0100-1.
- [26] and P. P. N. A. Tun, T. Chittin, N. Agarwal, M. L. New, Y. Thauang, "Tobacco Use among Young Adolescents in Myanmar: Findings from Global Youth Tobacco Survey," *Indian J. Public Health*, vol. 61, pp. 54–59, 2017, doi: 10.4103/ijph.IJPH.
- [27] B. Mbongwe, R. Taperia, N. Phaladze, A. Lord, and N. M. Zetola, "Predictors of smoking among primary and secondary school students in Botswana," *PLoS ONE*, vol. 12, no. 4, 2017, doi: 10.1371/journal.pone.0175640.
- [28] G. Lalithambigai, A. Rao, G. Rajesh, S. Ramya, and B. H. Mithun Pai, "Predictors of cigarette smoking among young adults in Mangalore, India," *Asian Pacific Journal of Cancer Prevention*, vol. 17, no. 1, pp. 45–50, 2016, doi: 10.7314/APJCP.2016.17.1.45.





- [29] L. Zhao, K. M. Palipudi, N. Ramanandraibe, and S. Asma, "Cigarette smoking and cigarette marketing exposure among students in selected African countries: Findings from the Global Youth Tobacco Survey," *Preventive Medicine*, vol. 91, pp. S35–S39, 2016, doi: 10.1016/j.ypmed.2015.12.015.
- [30] M. Maretalinia, E. Juliansyah, S. Suyitno, A. Yulianto, and D. Suryani, "Association of Smoking Related to Knowledge, Attitude, and Practice (KAP) with Tobacco Use in Community Health Center Working Area of Sungai Durian, Sintang Regency, West Kalimantan Province," *Buletin Penelitian Sistem Kesehatan*, vol. 24, no. 1, pp. 38–45, 2021, doi: 10.22435/hsr.v24i1.3544.
- [31] A. B. Barker, K. Whittamore, J. Britton, and J. Cranwell, "Content analysis of tobacco content in UK television," *Tobacco Control*, vol. 28, no. 4, pp. 381–385, 2019, doi: 10.1136/tobaccocontrol-2018-054427.
- [32] World Health Organization, "Banning tobacco advertising, promotion and sponsorship What you need to know," *World health Organization*, p. 13, 2013.
- [33] H. Blencowe *et al.*, "National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: A systematic analysis," *The Lancet Global Health*, vol. 4, no. 2, pp. e98–e108, 2016, doi: 10.1016/S2214-109X(15)00275-2.

BIOGRAPHIES OF AUTHORS







Sindu Setia Lucia     is a lecturer at Faculty of Health Science, Program of Administration and Health Policy, Respati Indonesia University, Yogyakarta, Indonesia. She can be contacted at email: sindulucia@gmail.com.







Novin Yetiani     is a lecturer in Midwifery Program, at Kapuas Raya College of Health Science, Sintang, West Kalimantan Province, Indonesia. Her study background is nursing and she has a lot of experiences of working in the hospital. She can be contacted at email: novin.yetiani@gmail.com.






Linda Suwarni     is a lecturer in Faculty of Health Science, Universitas Muhammadiyah Pontianak, Indonesia. She actively works in the field study of public health. She can be contacted at email: linda.suwarni@unmuhpnk.ac.id.






Heni Rusmitasari     is a lecturer at Faculty of Public Health, University of Muhammadiyah Semarang, Indonesia. She actively works as a lecturer and researcher in the area of interest administration and health policy. She can be contacted at email: heni.rusmitasari@unimus.ac.id.



Maretalinia Maretalinia    is a Ph. D student in Demography, Institute for Population and Social Research, Mahidol University, Thailand. She graduated from Bachelor of Public Health, Sriwijaya University and Master of Art (in Population and Sexual and Reproductive Health) Mahidol University. She used to work for two years as team-based Nusantara Sehat Batch II in Primary Health Care of Data Dian, Border Area of Indonesia-Malaysia. Her research of interest is maternal health and sociocultural of health. She can be contacted at email: mareta.linia.21@gmail.com.



Suyitno Suyitno    is a Master of Primary Healthcare Management. He graduated from ASEAN Institute for Health Development, Mahidol University, Thailand. He got the Bachelor of Public Health from University of Ahmad Dahlan, Yogyakarta, Indonesia. He is a research assistant at Indonesian Ministry of Health (Health Research and Development Agency). His field of research is environmental health and health development. He can be contacted at email: senopalawija@gmail.com.