



Linda Suwarni <linda.suwarni@unmuhpnk.ac.id>

[Global Health Management Journal] Result of Paper No.965: Major Revision (Due date: 13 November 2023)

2 pesan

Global Health Management Journal - GHMJ <ghmj.inschool@gmail.com>

6 November 2023 pukul 19.39

Kepada: linda.suwarni@unmuhpnk.ac.id

Cc: Doni <doni@inschool.id>, Global Health Management Journal - GHMJ <GHMJ@inschool.id>

Dear Dr. Linda Suwarni,

Regarding your submission with Paper No. 965 "Risk factors for severe stunted among Children aged 2-5 years with stunting in Pontianak City, Indonesia", the Editorial Boards of [Global Health Management Journal](#) are pleased to include it for next issue, after Major Revision.

Please find the attached file for your reference.

After revision, please send your Final Manuscript together with the result of Plagiarism Check no later than Monday, 13 November 2023.

During running the checking, please exclude the reference part. It is recommended to have less than 15% of similarity index; each reference similarity should not exceed 1%.

Thank you for trusting [Global Health Management Journal](#) for your home of publication.

Sincerely,

[Doni Marisi Sinaga](#)

Managing Editor

[Global Health Management Journal](#)

Yayasan Aliansi Cendekiawan Indonesia Thailand

Connect with us: [Facebook](#); [Instagram](#)[www.INSCHOOL.id](#); [Doni@inschool.id](#)**965-Article - Review Result.docx**

174K

Global Health Management Journal - GHMJ <ghmj.inschool@gmail.com>

22 November 2023 pukul 11.44

Kepada: linda.suwarni@unmuhpnk.ac.id

Cc: Doni <doni@inschool.id>, Global Health Management Journal - GHMJ <GHMJ@inschool.id>,

selviana@unmuhpnk.ac.id, Vidyastuti@unmuhpnk.ac.id, asrul.abdullah@unmuhpnk.ac.id, pranowo2054@gmail.com

Dear Dr. Linda Suwarni, [et.al.](#),**Congratulations!**

We are pleased to accept your submission "Risk factors for severe stunted among Children aged 2-5 years with stunting in Pontianak City, Indonesia" to be included in our incoming issue [Global Health Management Journal](#) Vol.6, No.2, 2023.

The online document is already available on our Website, or kindly visit the link below:

<https://publications.inschool.id/index.php/ghmj/article/view/965>

Attached in this email, the Letter of Acceptance from our Editor-in-Chief, for your reference.

We are very grateful for your contribution to the success of [Global Health Management Journal](#) in bringing our beneficiaries to a new world where knowledge and experiences can be shared and expanded.

Please feel free to reach me if any assistance is needed.

Best regards,

[Doni Marisi Sinaga](#)

Managing Editor

Global Health Management Journal

Yayasan Aliansi Cendekiawan Indonesia Thailand

Connect with us: [Facebook](#); [Instagram](#)


www.INSCHOOL.id; Doni@inschool.id

[Kutipan teks disembunyikan]



024. Linda Suwarni, et.al..pdf

153K



Open Access

Research Article

Risk factors for severe stunted among Children aged 2-5 years with stunting in Pontianak City, Indonesia

Linda Suwarni^{1*}, Selviana¹, Vidyastuti¹, Asrul Abdullah¹, Pranowo Adi²

¹ Fakultas Ilmu Kesehatan, Universitas Muhammadiyah Pontianak, Kalimantan Barat, Indonesia

² Widyaiswara Ahli Madya, BKKBN, Kalimantan Barat, Indonesia

*Corresponding author's email: linda.suwarni@unmuhpnk.ac.id

DOI: [10.35898/ghmj-XXXX](https://doi.org/10.35898/ghmj-XXXX)

© Yayasan Aliansi Cendekiawan Indonesia Thailand (Indonesian Scholars' Alliance). This is an open-access following Creative Commons License Deed - Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)

ABSTRACT

Background: Stunting is still a major public health in developing countries, including Indonesia. There are many predictors that might contribute to stunting, including child factors, mother factors, household factors, and community. This study focuses on children and mother level. **Objective:** This study aimed to examine the factors associated with severe stunted among children aged 2 to 5 years old. **Method:** This study uses primary data in Pontianak City, Indonesia the data has been collected from January to February 2023. Respondents were selected by total sampling method. Univariate, bivariate, and multivariate have been done using STATA 17. **Result:** The analysis data revealed that 75.98% of children were stunted and 24.20 were severe stunted. The factors including low birth weight and birth interval were found significantly associated with severe stunted, other independent variables did not have a correlation for being severe stunted. **Conclusion:** According to children's factors and maternal factors, the variables of low birth weight and birth interval were found to correlate with being severe stunted.

Keywords: Stunted, severe stunted, children, Pontianak

Received: 03 October 2022 **Reviewed:** 24 October 2022 **Revised:** 10 November 2022 **Accepted:** 25 November 2022

INTRODUCTION

Stunting among children under five years old is a significant public health issue globally, including in Indonesia and Asian countries (Correa, 2022; Mengesha et al., 2021). Stunting is defined as a condition where a child's height or length is significantly below the average for their age (De Onis & Branca, 2016). It is estimated that nearly one-third of children under five years old worldwide are affected by stunting, with higher prevalence in low-resource countries in sub-Saharan Africa and South Asia (Mengesha et al., 2021). Stunting among children under five years old is a significant public health issue in Indonesia and Asian countries (Umar & Darajat, 2022). Stunting is defined as a child's length or height being below the acceptable standard values for their age (De Onis & Branca, 2016). It is estimated that nearly one-third of children under five years old worldwide are affected by stunting, with higher prevalence in low-resource countries in sub-Saharan Africa and South Asia (Mengesha et al., 2021). In Indonesia, 30.8% of children under five

Commented [Office1]: Introduction should provide information or rough details of the keyword at your title, including **Stunting Risk Factors**, why it is important to study about that? Start by saying the health benefits and the consequences if neglecting this issue.

Find the theory related! Show if there are any examples/applications in this world?

Raise the National Issue then continue explaining the urgent to conduct this research for the sampling locations.

Why **Pontianak**? Any incidence? Or the results of this study can be used for local regulation? Show to the reader why this study is important.

Then, authors can show the urgency of this research. At that moment, the authors can clearly state the aims of this study at the end of the section.

experience stunting (Umar & Darajat, 2022). Factors associated with stunting can be categorized into several clusters, including maternal nutrition and infection, teenage motherhood and short birth intervals, fetal growth restriction and preterm birth, child nutrition and infection, and environmental factors (Danaei et al., 2016).

Stunting, defined as low height-for-age, is a significant public health issue affecting children under 5 years old worldwide. It is associated with long-term consequences such as poor cognitive development, increased risk of chronic diseases, and reduced economic productivity (Lartey, 2015). Understanding the factors related to severe stunting is crucial for developing effective interventions to address this problem. Several studies have investigated these factors in different countries, providing valuable insights into the determinants of severe stunting among children under 5 years old. One study conducted in Nigeria found that factors such as maternal education, household wealth, and access to improved drinking water sources were significantly associated with severe stunting among children under 5 years old (Akombi et al., 2017). Similarly, a study in Rwanda identified risk factors for stunting, including household wealth, maternal education, and access to improved sanitation facilities (Nshimiyiryo et al., 2019). These findings highlight the importance of socioeconomic factors in determining the prevalence of severe stunting. Another study conducted in Indonesia found that maternal education, maternal occupation, age at pregnancy, socioeconomic conditions, and nutritional status of pregnant women were associated with stunting among children under 2 years old (Susyani et al., 2022). This suggests that improving maternal education and socioeconomic conditions can play a crucial role in preventing severe stunting in early childhood.

In Nepal, a study found that factors such as delivery complications, maternal education, household wealth, and access to healthcare services were associated with stunting and severe stunting among children under 5 years old (Tiwari et al., 2014). This highlights the importance of addressing both maternal and environmental factors in preventing severe stunting. The role of maternal factors in stunting was also highlighted in a study conducted in Ghana, which found that young maternal age was a risk factor for child undernutrition (Wemakor et al., 2018). Similarly, a study in Ethiopia reported that the quantity, frequency, and type of supplementary feeding, birth weight, sex, birth order, and disease conditions like diarrhea were strongly associated with stunting among under-five children (Fikadu et al., 2014). Furthermore, studies conducted in Zambia and Bangladesh identified factors such as sex and age of the child, mother's age and level of education, wealth status, improved source of drinking water, duration of breastfeeding, and residence as predictors of stunting among children under 5 years old (Chowdhury et al., 2021; Mzumara et al., 2018). These findings emphasize the importance of addressing multiple factors at the individual, household, and community levels to prevent severe stunting.

In addition to socioeconomic and maternal factors, other studies have identified specific risk factors for severe stunting. For example, a study in Ethiopia found that children from educated and malnourished mothers, as well as those from less wealthy mothers, were more likely to be severely stunted (Hailu et al., 2020). Another study in Rwanda reported that male gender, age above 11 months, child disability, having two children below the age of five, and a child having diarrhea were associated with increased odds of childhood stunting (Kalinda et al., 2023). Environmental factors, such as sanitation and access to clean water sources, have also been identified as important determinants of severe stunting. A study in Zimbabwe highlighted the low micronutrient density and poor protein quality in cereal-based diets as contributing factors to stunting in resource-poor settings (Kairiza et al., 2020). Similarly, a study in Bangladesh found that toilet facilities, source of drinking water, place of residence, and socioeconomic status were significantly associated with stunting, wasting, and being underweight in children under five (Chowdhury et al., 2021).

Overall, the factors related to severe stunting among children under 5 years old are multifaceted and include socioeconomic factors, maternal factors, environmental factors, and specific risk factors. Addressing these factors requires a comprehensive approach that includes improving maternal education, socioeconomic conditions, access to healthcare services, and environmental sanitation. Additionally, interventions should focus on promoting optimal breastfeeding practices, improving the quality and diversity of children's diets, and providing access to clean water sources. By addressing these factors, it is possible to reduce the prevalence of severe stunting and improve the health and well-being of children under 5 years old.

METHODOLOGY

This study used a cross-sectional time frame. The data collection has been done in 10 villages in 6 sub-districts in Pontianak City, West Kalimantan Province where stunting is the focus. Those locations such as Parit Tokaya, Benua Melayu Laut, Tambelan Sampit, Banjar Serasan, Saigon, Tanjung Hilir,

Commented [Office2]: Attention to following guidelines/subsections, and complete it at your methods

* **Study design/ Research procedures:** Type of research? Sample size? Who is the key informant? How you select them? Sampling method? From where, and who are they? Provide inclusion and exclusion criteria! Any are drop outs? how you get the samples?

* **Measurements**

– Show the independent and dependent variable here! How you get the data? Questionnaire? Observation? Mention and describe the steps clearly!

– Preparation (...)?

– Who did the measurement? If the authors, please state in detail the way to measure the parameters! If researcher from other institution, please mention the name of organization, methods/procedures, lab accreditation (if working in lab)!

– Elaborate how you define the level! Scoring? What questions? References?

* **Statistical techniques** applied – what analysis you run? Also provide the reason/purposes of each tests!

* **Ethical Clearance:** Institution/Provider, Number of Letter, Date issued, any consent offered before involved their participation?

Tanjung Hulu, Parit Mayor, Siantan Hulu dan Sungai Jawi. The data was collected during January to February 2023. Total sampling was used to select the respondents who were mothers of children aged 2 to 5 years old with diagnosed stunting. The final respondents who completely joined the interview were 687 mothers with children aged 2-5 years old.

The dependent variable of this study is nutritional status which has two categories, such as stunted and severe stunted. To define severe stunted, authors used cut-off based on WHO recommendation Length/height-for-age <-3 SD of the median, and stunted used Length/height-for-age ≤-2 SD and ≥-3 SD of the median. The independent variable of this study includes the children's level and mother level.

Data analysis in this study includes univariate, bivariate, and multivariate. Univariate was used to describe the general information of respondents that presented in frequency and percentage. Bivariate analysis was done using Chi-square in order to test each independent variable and dependent variable. Multivariate analysis in this study is binary logistic regression because dependent variable consists of dummy categories. All the analysis was done by using STATA 17. This study including instruments and all procedure has been approved by Ethical Clearance No: 010/KEPK-FIKES/UM PONTIANAK/2023.

RESULTS

Univariate Analysis

Univariate analysis in this study explains the frequency distribution of children with stunting characteristics and the potential predictors. In Table 1. Among all stunting children, there are 165 of 687 children who were severely stunted (24.20%). According to the birth interval between childbirth and the previous birth, it was revealed that the majority of children were born in intervals of 2 years or more with another birth (73.80%). Moreover, the gestation data showed the majority of children were born at enough months of gestation (93.16%). The mother reported that the children were born with enough or more birth weight (83.84%) and enough birth height (55.75%). Based on the mother's history of chronic lack of energy, the mother reported did not have (87.34%). The majority of mothers practiced exclusive breastfeeding (80.35%). Furthermore, it also described that most of the mothers had a weight of 150 cm or higher (66.81%), did not have a history of anemia (65.36%), and did not complete all immunizations (56.62%).

Table 1. The General Information of Children

Variables (n=687)	Frequency	Percentage
Stunting		
Severely Stunted	165	24.20
Stunted	522	75.98
Birth interval		
< 2 years	180	26.20
\geq 2 years	507	73.80
Premature when born		
Yes	47	6.84
No	640	93.16
Low birth weight		
Yes	111	16.16
No	576	83.84
Birth height		
< 48 cm	304	44.25
\geq 48 cm	383	55.75
History of chronic lack of energy		
Yes	87	12.66
No	600	87.34
Exclusive Breastfeeding		
Yes	136	19.80
No	551	80.20
Age when pregnant (year)		

Commented [Office3]: Questionnaire? Open questions? Please make it details. Who and how to measure the variables?

Commented [Office4]: For the title of sub section, try to not use Univariate and soon, but something easy to read, and can represent the content of that subsection.

< 20 th or >35 th	552	80.35
20-35	135	19.65
Mother's height		
< 150 cm	228	33.19
≥ 150 cm	459	66.81
Mother's History of Anemia		
Yes	238	34.64
No	449	65.36
Complete all immunization		
No	389	56.62
Yes	298	43.38
Total	687	100.0

Bivariate Analysis

Table 2 below shows the result of the Chi-square test by bringing each potential predictor associated with severe stunted. It showed that some variables including birth interval, premature birth, low birth weight, and birth height have strong associations ($p\text{-value} < 0.001$) with severe stunted. Moreover, another variable has a moderate association ($p\text{-value} < 0.01$), including exclusive breastfeeding, and a low association ($p\text{-value} < 0.05$) including the history of chronic lack of energy. However, some other variables were not significantly associated with nutritional status ($p\text{-value} > 0.05$), including age when pregnant, mother's height, history of anemia, and completeness of immunizations.

Table 2. The result of bivariate analysis using the Chi-square test

Variables	Nutritional status		<i>p-value</i>
	Severe stunted (%)	Stunted (%)	
Birth interval			< 0.001***
< 2 years	61 (33.89)	119 (66.11)	
≥ 2 years	104 (20.51)	403 (79.49)	
Premature when born			< 0.001***
Yes	27 (57.45)	20 (42.55)	
No	138 (21.56)	502 (78.44)	
Low birth weight			< 0.001***
Yes	62 (55.86)	49 (44.14)	
No	103 (17.88)	473 (82.12)	
Birth height			< 0.001***
< 48 cm	95 (31.25)	209 (68.75)	
≥ 48 cm	70 (18.28)	313 (81.72)	
History of chronic lack of energy			0.014*
Yes	30 (34.48)	57 (65.52)	
No	135 (22.50)	465 (77.50)	
Exclusive Breastfeeding			0.006**
Yes	45 (33.09)	91 (66.91)	
No	120 (21.78)	431 (78.22)	
Age when pregnant (year)			0.586
< 20 th or >35 th	135 (24.46)	417 (75.54)	
20-35	30 (22.22)	105 (77.78)	
Mother's height			0.814
< 150 cm	56 (24.56)	172 (75.44)	
≥ 150 cm	109 (23.75)	350 (76.25)	
Mother's History of Anemia			0.730
Yes	59 (24.79)	179 (75.21)	
No	106 (23.61)	343 (76.39)	
Complete all immunization			0.085
No	103 (26.48)	286 (73.52)	
Yes	62 (20.81)	236 (79.19)	

* $p\text{-value} < 0.05$, ** $p\text{-value} < 0.001$, *** $p\text{-value} < 0.001$

Multivariate Analysis

Table 3 below describes the result of multivariate analysis using binary logistic regression. It was shown that, after adjusting to all independent variables, compared to children with enough birth weight, the variables of having low birth weight were 4.26 times more likely and strongly associated with severe stunted (p value = ...). Additionally, compared to children born with 2 year or more birth interval, children born with less than a 2-year birth interval had a 45% increasing probability of having severe stunted. However, after adjusting to all predictors the variables of premature when born, birth height, history of chronic lack of energy, exclusive breastfeeding, age when pregnant, mother's height, mother's history of anemia, and completeness of immunizations were found not significantly associated with having severe stunted (p -value>0.05).

Table 3. The result of multivariate analysis using binary logistic regression

Variables	AOR	95% Confidence Interval (Lower – Upper)	<i>p</i> -value
Birth interval (ref: ≥ 2 years)			
< 2 years	0.55	0.37 – 0.81	0.002**
Premature when born (ref: No)			
Yes	1.91	0.93 – 3.92	0.080
Low birth weight (ref: No)			
Yes	4.26	2.55 – 7.13	0.000***
Birth height (ref: ≥ 48 cm)			
< 48 cm	1.22	0.81 – 1.84	0.334
History of chronic lack of energy (ref: No)			
Yes	1.64	0.95 – 2.84	0.075
Exclusive Breastfeeding (ref: Yes)			
No	1.51	0.96 – 2.38	0.074
Age when pregnant (year) (ref: 20-35)			
< 20 th or >35 th	0.68	0.41 – 1.21	0.131
Mother's height (ref: ≥ 150 cm)			
< 150 cm	0.94	0.63 – 1.41	0.757
Mother's History of Anemia (ref: No)			
Yes	0.94	0.63 – 1.41	0.757
Complete all immunizations (ref: Yes)			
No	1.33	0.91 – 1.97	0.143

* p -value <0.05, ** p -value <0.001, *** p -value <0.001

LR chi2 (10) = 85.73

Prob > chi2 = 0.000

Pseudo R2 = 0.1132

Log likelihood = -335.86

DISCUSSION

According to the result in the previous section, the factors associated with severe stunted was low birth weight and birth interval. This finding was in line with the studies in developing countries (Danaei et al., 2016). Several factors have been identified as being associated with stunting among children under five years old. These factors can be categorized into different clusters, including maternal nutrition and infection, teenage motherhood and short birth intervals, fetal growth restriction and preterm birth, child nutrition and infection, and environmental factors (Danaei et al., 2016). Maternal nutrition and infection play a crucial role in the development of stunting, as poor maternal nutrition and infections during pregnancy can lead to fetal growth restriction and preterm birth, which are risk factors for stunting (Danaei et al., 2016). Teenage motherhood and short birth intervals are also associated with stunting, as these factors can lead to inadequate maternal nutrition and care during pregnancy (Danaei et al., 2016). Child nutrition and infection are important factors in the development of stunting. Inadequate nutrition, including insufficient intake of essential nutrients, can contribute to stunting (Marume et al., 2023). Additionally, frequent infections, such as diarrheal and worm infections, can impair nutrient absorption and utilization,

leading to stunting (Marume et al., 2023). Environmental factors, such as access to safe water, sanitation, and hygiene practices, also play a role in stunting. Poor access to safe water and sanitation facilities can increase the risk of infections, which can contribute to stunting (Shoffifah et al., 2022). Other factors associated with stunting include low birth weight, low socioeconomic status, inadequate breastfeeding practices, and poor dietary quality (Marume et al., 2023). Low birth weight is a significant risk factor for stunting, as it reflects inadequate fetal growth and development (Marume et al., 2023). Socioeconomic factors, such as low income and low maternal education, are also associated with stunting, as they can limit access to nutritious food and healthcare (Marume et al., 2023). Inadequate breastfeeding practices, including early cessation of breastfeeding and lack of exclusive breastfeeding, can contribute to stunting (Marume et al., 2023). Poor dietary quality, including insufficient intake of essential nutrients, can also contribute to stunting (Marume et al., 2023).

Several child-related factors have been identified as predictors of stunting, including child's age, birth length, birth weight, weight-for-age outcome, breastfeeding status, number of meals, dietary quality, child's appetite, and presence of diarrheal and worm infections (Marume et al., 2023). Maternal factors such as age, education, occupation, and HIV status also play a role in stunting (Marume et al., 2023). Household and socio-cultural factors, such as income status, access to safe water and toilets, health clubs, and maternal support in infant feeding, are also significant predictors of childhood stunting (Marume et al., 2023). Other studies have identified additional risk factors for stunting, including low paternal education, maternal height less than 150 cm, high-risk maternal age, low birth weight, and history of infectious diseases (Fikadu et al., 2014; Manggala et al., 2018). The consumption of animal-sourced foods, child underweight status, and income type have also been associated with stunting (Umwali et al., 2022). It is important to note that stunting has long-term consequences for children's health, well-being, and productivity (Correa, 2022). Stunted children are more susceptible to infections and are at a higher risk of cognitive, motor, and language impairments (Park et al., 2019). Stunting also has economic implications, as it can lead to decreased productivity and increased risk of chronic diseases in adulthood (McGovern et al., 2017; Nasser et al., 2022).

It is important to note that stunting is a complex issue influenced by multiple factors, and these factors can interact with each other. For example, maternal nutrition and infection can affect fetal growth and development, which can then impact child nutrition and infection (Danaei et al., 2016). Additionally, socioeconomic factors can influence access to nutritious food and healthcare, which can in turn affect child nutrition and infection (Marume et al., 2023). Efforts to reduce stunting should focus on improving maternal nutrition and infection control, promoting optimal child nutrition and reducing the burden of childhood illnesses, and addressing environmental factors such as access to safe water and sanitation (Danaei et al., 2016). Additionally, interventions targeting household and socio-cultural factors, such as improving income status and maternal support in infant feeding, can also contribute to reducing stunting (Marume et al., 2023).

In conclusion, stunting among children under five years old in Indonesia and Asian countries is a complex issue influenced by various factors. These factors include maternal nutrition and infection, teenage motherhood and short birth intervals, fetal growth restriction, and preterm birth, child nutrition and infection, environmental factors, low birth weight, socioeconomic status, breastfeeding practices, and dietary quality. Efforts to address stunting should focus on improving maternal nutrition and healthcare, promoting exclusive breastfeeding, improving access to nutritious food, implementing effective water, sanitation, and hygiene practices, and addressing socioeconomic factors. By addressing these factors comprehensively, it is possible to reduce the prevalence of stunting and improve the health and well-being of children in Indonesia and Asian countries. In conclusion, stunting among children under five years old is a significant public health issue in Indonesia and Asian countries. It is influenced by a range of factors, including child-related, maternal, household, and socio-cultural factors. Efforts to reduce stunting should focus on addressing these factors through interventions that improve maternal and child nutrition, reduce

Commented [Office5]: Mention again here about the result of your study in Pontianak!

the burden of childhood illnesses, and improve access to safe water and sanitation. These interventions have the potential to improve the health, well-being, and productivity of children in these countries.

CONCLUSION

Stunting is a major public health issue in developing countries, including Indonesia. In high-risk areas of stunting in Pontianak City, Indonesia it was found 75.98% of children with a high risk of stunting were severe stunted, and 24.20% were stunted. According to children's factors and maternal factors, the variables of low birth weight and birth interval were found to correlate with being severe stunted.

These findings may lead to policy recommendations to encourage women to have at least two years of birth interval and increase intake of nutritious food during pregnancy to prevent low birth weight. All other stakeholders also need to promote and educate women and family members about the prevention of stunting starting from pregnancy period.

Commented [Office6]: Your results are tremendous, try to cover all of that in the Conclusion part. Not only cover the significant factors for being severe stunted, but also mention the others factors (Only mention, no need to elaborate).

Commented [Office7]: Nice wrap up!

REFERENCES

- Akombi, B. J., Agho, K. E., Hall, J. J., Merom, D., Astell-Burt, T., & Renzaho, A. M. N. (2017). Stunting and severe stunting among children under-5 years in Nigeria: A multilevel analysis. *BMC Pediatrics*, *17*(1), 15. <https://doi.org/10.1186/s12887-016-0770-z>
- Chowdhury, M. R. K., Khan, H. T. A., Mondal, Md. N. I., & Kabir, R. (2021). Socio-demographic risk factors for severe malnutrition in children aged under five among various birth cohorts in Bangladesh. *Journal of Biosocial Science*, *53*(4), 590–605. <https://doi.org/DOI: 10.1017/S0021932020000425>
- Correa, P. R. (2022). Factors associated with stunting among children 0 to 59 months of age in Angola: A cross-sectional study using the 2015–2016 Demographic and Health Survey. *PLOS Global Public Health*, *2*(12), e0000983.
- Danaei, G., Andrews, K. G., Sudfeld, C. R., Fink, G., McCoy, D. C., Peet, E., Sania, A., Smith Fawzi, M. C., Ezzati, M., & Fawzi, W. W. (2016). Risk factors for childhood stunting in 137 developing countries: a comparative risk assessment analysis at global, regional, and country levels. *PLoS Medicine*, *13*(11), e1002164.
- De Onis, M., & Branca, F. (2016). Childhood stunting: a global perspective. *Maternal & Child Nutrition*, *12*, 12–26.
- Fikadu, T., Assegid, S., & Dube, L. (2014). Factors associated with stunting among children of age 24 to 59 months in Meskan district, Gurage Zone, South Ethiopia: a case-control study. *Bmc Public Health*, *14*(1), 1–7.
- Hailu, B. A., Bogale, G. G., & Beyene, J. (2020). Spatial heterogeneity and factors influencing stunting and severe stunting among under-5 children in Ethiopia: spatial and multilevel analysis. *Scientific Reports*, *10*(1), 16427. <https://doi.org/10.1038/s41598-020-73572-5>
- Kairiza, T., Kembo, G., Pallegedara, A., & Macheke, L. (2020). The impact of food fortification on stunting in Zimbabwe: does gender of the household head matter? *Nutrition Journal*, *19*(1), 22. <https://doi.org/10.1186/s12937-020-00541-z>
- Kalinda, C., Phiri, M., Qambayot, M. A., Ishimwe, M. C. S., Gebremariam, A., Bekele, A., & Wong, R. (2023). Socio-demographic and environmental determinants of under-5 stunting in Rwanda: Evidence from a multisectoral study. *Frontiers in Public Health*, *11*, 1107300.
- Lartey, A. (2015). What would it take to prevent stunted growth in children in sub-Saharan Africa? *Proceedings of the Nutrition Society*, *74*(4), 449–453. <https://doi.org/DOI: 10.1017/S0029665115001688>
- Manggala, A., Kenwa, K. W., Kenwa, M., Sakti, A. A. G. D. P., & Sawitri, A. A. (2018). Risk factors of stunting in children aged 24-59 months. *Paediatrica Indonesiana*, *58*(5). <https://doi.org/10.14238/pi58.5.2018.205-12>
- Marume, A., Archary, M., & Mahomed, S. (2023). Predictors of stunting among children aged 6–59 months, Zimbabwe. *Public Health Nutrition*, *26*(4), 820–833. <https://doi.org/DOI: 10.1017/S1368980023000046>
- McGovern, M. E., Krishna, A., Aguayo, V. M., & Subramanian, S. V. (2017). A review of the evidence linking child stunting to economic outcomes. *International Journal of Epidemiology*, *46*(4), 1171–1191. <https://doi.org/10.1093/ije/dyx017>

- Mengesha, A., Hailu, S., Birhane, M., & Belay, M. M. (2021). The Prevalence of Stunting and Associated Factors among Children Under Five years of age in Southern Ethiopia: Community Based Cross-Sectional Study. *Annals of Global Health*. <https://doi.org/10.5334/aogh.3432>
- Mzumara, B., Bwembya, P., Halwiindi, H., Mugode, R., & Banda, J. (2018). Factors associated with stunting among children below five years of age in Zambia: evidence from the 2014 Zambia demographic and health survey. *BMC Nutrition*, 4(1), 51. <https://doi.org/10.1186/s40795-018-0260-9>
- Nasser, M. S., Akhlaq, A., Ali, H., & Nasser, D. (2022). Evidence Linking Stunting to Economic Outcomes in the SAARC Region: A Systematic Review. *Pakistan Journal of Public Health*, 12(2), 88–101.
- Park, J. J. H., Harari, O., Siden, E., Dron, L., Zannat, N.-E., Singer, J., Lester, R. T., Thorlund, K., & Mills, E. J. (2019). Interventions to improve linear growth during complementary feeding period for children aged 6-24 months living in low-and middle-income countries: a systematic review and network meta-analysis. *Gates Open Research*, 3.
- Shofifah, A., Sulistyorini, L., & Praveena, S. M. (2022). Environmental Sanitation at Home and History of Infection Diseases as risk Factors for Stunting in Toddlers in Drokilo Village, Kedungadem District, Bojonegoro Regency. *Journal of Environmental Health*, 14(4).
- Susyani, S., Febry, F., Margareth, I., Sadiq, A., Sartono, S., Sari, I. P., & Ni'mah, T. (2022). Maternal Risk Factor on Incidence of Stunting in South Sumatera. *Open Access Macedonian Journal of Medical Sciences*, 10(E), 1599–1604.
- Tiwari, R., Ausman, L. M., & Agho, K. E. (2014). Determinants of stunting and severe stunting among under-fives: evidence from the 2011 Nepal Demographic and Health Survey. *BMC Pediatrics*, 14(1), 239. <https://doi.org/10.1186/1471-2431-14-239>
- Umar, E., & Darajat, A. (2022). Factors Affecting the Management of Stunting in Banten Province. *KnE Life Sciences*, 380–393.
- Umwali, N., Kunyanga, C. N., & Kaindi, D. W. M. (2022). Determinants of stunting in children aged between 6–23 months in Musanze region, Rwanda. *Frontiers in Nutrition*, 9. <https://www.frontiersin.org/articles/10.3389/fnut.2022.1044350>
- Wemakor, A., Garti, H., Azongo, T., Garti, H., & Atosona, A. (2018). Young maternal age is a risk factor for child undernutrition in Tamale Metropolis, Ghana. *BMC Research Notes*, 11(1), 877. <https://doi.org/10.1186/s13104-018-3980-7>

Cite this article as:

Suwarni, Linda. Risk factors for severe stunted among Children aged 2-5 years with stunting in Pontianak City, Indonesia. *GHMJ (Global Health Management Journal)*. 2023; 5(1):01-05. doi:10.35898/ghmj-XXXX